

Adaptive Uncertainty Resolution in Bayesian Combinatorial Optimization Problems*

Sudipto Guha
University of Pennsylvania

Kamesh Munagala
Duke University

January 28, 2010

Abstract

In several applications such as databases, planning, and sensor networks, parameters such as selectivity, load, or sensed values are known only with some associated uncertainty. The performance of such a system (as captured by some objective function over the parameters) is significantly improved if some of these parameters can be probed or observed. In a resource constrained situation, deciding which parameters to observe in order to optimize system performance, itself becomes an interesting and important optimization problem. This general problem is the focus of this paper.

One of the most important considerations in this framework is whether adaptivity is required for the observations. Adaptive observations introduce blocking or sequential operations in the system whereas non-adaptive observations can be performed in parallel. One of the important questions in this regard is to characterize the benefit of adaptivity for probes and observation.

We present general techniques for designing constant factor approximations to the optimal observation schemes for several widely used scheduling and metric objective functions. We show a unifying technique that relates this optimization problem to the *outlier* version of the corresponding deterministic optimization. By making this connection, our technique shows constant factor upper bounds for the benefit of adaptivity of the observation schemes. We show that while probing yields significant improvement in the objective function, being adaptive about the probing is not beneficial beyond constant factors.

1 Introduction

Consider a measurement scenario such as sensor networks or spectrograms, where we have errors in estimation, but the distribution of the error is understood through models of the measuring instrument, historical data, regression, Kalman filters, etc. For example, suppose we have a set of data-points and are trying to construct a classification of them using the minimum spanning tree (MST). Suppose that we cannot resolve all the errors but we can repeat and refine a few of the measurements, i.e., “probe” a few points to make detailed measurements. A natural question

*Parts of this paper appeared in the Proceedings of the 18th ACM-SIAM Symposium on Discrete Algorithms, 2007 [18].

Author’s address: Sudipto Guha, Department of Computer and Information Sciences, University of Pennsylvania. Supported in part by an Alfred P. Sloan Research Fellowship, by NSF awards CCF-0644119, and CNS-0721541. Email: sudipto@cis.upenn.edu.

Kamesh Munagala, Department of Computer Science, Duke University, Durham NC 27708. Research supported by an Alfred P. Sloan Research Fellowship, and by NSF via a CAREER award and grant CNS 0540347. Email: kamesh@cs.duke.edu.

is which measurements do we refine, say if we wanted to refine only k of them and wanted to optimize the expected cost of the MST? In many situations, we can assume that the errors made by different instruments are independent of each other – even then the problem is NP-HARD [15] for computing very simple functions such as the *minimum* of the readings. In this paper we focus on general combinatorial optimization problems.

The above is a typical example of a model driven optimization problem [15]. This has gained significant currency in a variety of other research areas such as database query optimization and route selection in networks. In a database query optimization setting suppose the optimizer is presented with a set of sufficiently complicated and unrelated (independent) queries. The query-optimizer can estimate the resources needed by the queries from historical information, cached statistics, sampling of various sub-queries, or by performing inexpensive filters [3, 8, 9]. Subsequent to this estimation, the query optimizer schedules the tasks to optimize the throughput or the average completion times. However, this estimation process itself consumes resources such as time, network bandwidth, and space, and therefore the decision to choose the parameters to refine in estimate becomes a key optimization problem. A similar problem arises in networking, where the current state of multiple routes and servers can be probed and observed before deciding which route/server to use for a specific connection [1, 21, 20]. The above examples can also be extended to planning over a network, where we may wish to visit a sequence of nodes, about which we have imprecise information and can only refine a small number of them, to minimize total distance traveled – such problems arise in query processing in sensor networks [12].

Problem Statement. The above problems of measurement refinement or optimizing independent query schedules can be formulated abstractly as follows. Let $X_1, X_2, \dots, X_n$ be non-negative independent random variables, whose distributions are given as inputs (the distribution can be specified by samples and does not affect any result in this paper). Let $[n]$ denote $\{1, 2, \dots, n\}$. For subset $S \subseteq [n]$, let $\mathbf{v}_S$ denote the realization of $\{X_i, i \in S\}$. We are given a minimization problem $h(o; \mathbf{v})$, where $\mathbf{v} \in \mathbb{R}_+^n$ denotes the inputs, and $o \in \mathcal{O}$ denotes the output.

As an example, in weighted completion time scheduling on a single machine, $1||\sum_j w_j C_j$, job i has size that follows distribution X_i , and has weight w_i . The vector $\mathbf{v}$ denotes a realization of job sizes, the space $\mathcal{O}$ is the set of all orderings of jobs, and the function $h(o; \mathbf{v}) = \sum_i w_i \sum_{j \prec_o i} \mathbf{v}_j$, where $j \prec_o i$ means job j appears before i in the ordering o . Note that if the goal is to solve $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}}[h(o; \mathbf{v})]$, the optimal o simply orders the jobs by Smith's rule in decreasing order of $\frac{w_i}{\mathbf{E}[X_i]}$. If the goal is instead to solve $\mathbf{E}_{\mathbf{v}}[\min_{o \in \mathcal{O}} h(o; \mathbf{v})]$, for each realization $\mathbf{v}$, the ordering would be in decreasing $\frac{w_i}{\mathbf{v}_i}$. Our goal will not be to solve either of these extreme versions.

Our goal will be to solve the probing version of these problems. Here, each variable $i \in [n]$ has an observation cost c_i . The exact value of the underlying random variable X_i can be found by spending this cost. There is a budget $C (\geq \max_i c_i)$ on the total cost of observation. The probing policy chooses a subset of variables of total cost at most C to probe and observe. *Subsequent* to this, the optimization policy chooses a solution based on the outcome of the probes. We note that the solution is fixed *after* the outcomes of the probes, but *before* the realization of the unprobed variables. Therefore, given the set of probed variables and their outcomes, the solution is the same for *all* realizations of the unprobed variables. The final goal is to devise a policy (or strategy) for adaptively choosing that $S \subseteq [n]$ of variables so that $\mathbf{E}_{\mathbf{v}_S} [\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]]$ is minimized. The expectation is jointly over outcomes of probes and the distributions of unprobed variables. Here, $\mathbf{v}_S$ denotes the realization of the random variables $\{X_i, i \in S\}$, and $\bar{S}$ denotes $[n] \setminus S$.

In the above completion time example, let $\mathbf{E}[X_i] = \mu_i$. If the probing policy chooses a subset S of jobs to probe, after the probing, the optimal ordering is Smith's Rule using the exact sizes $\mathbf{v}_i$ for the probed jobs and the expected sizes μ_i of the unprobed jobs. For a given realization $\mathbf{v}_S$ of the

variables $\{X_i, i \in S\}$, the expected completion time (where the expectation is over the unprobed job sizes) is:

$$\begin{aligned} \min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})] &= \sum_{i,j \in S, j \geq i} \min(w_i \mathbf{v}_j, w_j \mathbf{v}_i) \\ &+ \sum_{i,j \notin S, j \geq i} \min(w_i \mu_j, w_j \mu_i) + \sum_{i \in S, j \notin S} \min(w_i \mu_j, w_j \mathbf{v}_i) \end{aligned}$$

The value of the overall solution given S is probed is therefore:

$$\mathbf{E}_{\mathbf{v}_S} \left[\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})] \right] \quad (1)$$

The objective of the probing strategy is to adaptively choose a subset S to probe, so that this value is minimized, subject to $\sum_{i \in S} c_i \leq C$. Note that such a strategy is a decision tree with the next decision to probe depending on the outcomes so far. At a leaf l of the decision tree, a subset S_l has been probed and realized; the job ordering at the leaf would have weighted completion time given by Equation (1) with $S = S_l$. The expected value of this quantity over the decision tree is the value of the probing strategy. The goal is to design a polynomial time algorithm for finding the optimal probing strategy.

Conceptual Issues. Several conceptual and systems issues arise immediately.

- The most important consideration for a system is quantifying the benefit of being adaptive in the probing policy. In the above formulation, the probes could possibly all be done in parallel upfront, *i.e.*, non-adaptively, and the results of the probes are ignored in deciding which other variables to probe. In contrast, adaptive observations are sequential, based on the outcomes of the previous probes. It is clear that adaptive strategies yield as good or better solutions than the non-adaptive counterpart. The first issue with an adaptive strategy is the complexity of expressing the solution; it is *a priori* not clear that the decision tree that encodes the optimum probing policy (or strategy) is polynomially bounded, so that it is not clear if the problem of finding the optimal adaptive probing policy is even in NP.
- The next interesting issue, which is a consequence of the adaptivity, is the budget. Note that while probing, if a suitable value is found already, then the probing can be halted early, yielding significant cost savings. Thus it makes sense to assume that the budget is in expectation as well. This sets up two classes of problems, namely with *hard budgets*, where the probing budget is C is on all decision paths, and with *soft budgets*, *i.e.*, the budget C is in expectation over the decision paths, so that on some paths, the cost of the variables probed could exceed C if this is compensated by halting early on other decision paths. Note that this distinction does not arise for non-adaptive policies, where the probes are done upfront in parallel.

Results. In this paper, we design poly-time algorithms for finding approximately optimal *non-adaptive* probing strategies for the large class of scheduling and metric objective functions that have been considered in the known applications and literature to date, namely, weighted completion time scheduling, minimum makespan scheduling, metric clustering, and Steiner trees. Our results for all these seemingly disparate problems use a common underlying technique that also shows that these strategies have a bounded *adaptivity gap*, *i.e.*, there exists a non-adaptive probing strategy which is only a constant factor worse compared to the best adaptive probing strategy with soft

budgets. Therefore for the problems we study, the non-adaptive and adaptive models with hard and soft budgets are all related to each other by constant factors. As a consequence, one of the main points we establish is that although probing helps significantly¹, adaptive probing is no better than non-adaptive probing by more than a constant factor in many problems of interest.

Related Work. The model-driven optimization framework can be defined for other objective functions h . This class of problems was first defined by the authors of [15], where the non-adaptive MINIMUM-ELEMENT problem, with $h = \min_i X_i$ is considered. Though the current paper presents the best results known for scheduling and metric problems, other work [15, 18, 19, 20] has considered different objective functions, most notably problems involving subset selection. In [18, 19], we considered the case where f is a single constraint packing problem such as knapsack with random profits which are observable, and present a 8 approximation based on rounding the solution of a natural linear program. As in this paper, the approximation ratio holds even when the optimal solution is allowed adaptive (*i.e.*, can be based on the results of previous observations). It further holds even when the hidden quantity is a distribution (instead of a single value) and a prior on this distribution is specified as input. In [20] the authors consider the Lagrangean version of $h = \max_i X_i$, where the observations are adaptive and the goal is to maximize the expected difference between the maximum value and the observation cost. Note that there is no budget on this cost, instead it is part of the objective function. We note that the techniques needed for subset selection problems are very different from those needed for the scheduling and metric problems considered in this paper, where all variables whether probed or unprobed eventually are part of the solution.

The notion of refining uncertainty has been considered in an *adversarial setting* by several researchers [31, 13, 27, 4]. Here, the only prior information about an input is the lower and upper bounds on its value. The goal is to minimize the observations needed to estimate some function over these inputs *exactly*, and often strong lower bounds arise. For correlated random variables, the problem of minimizing residual information is considered in [29].

The adaptivity gap in the absence of probes has been considered in the literature earlier, notably in [10, 11, 30, 35] for Knapsack and scheduling problems. However, the model driven optimization problem is considerably different from the settings in those papers. In [10, 11, 30, 35] the optimum is allowed to decide on the next item to schedule based on the past, but once the next item is decided this is an irrevocable commitment. In contrast, in our problem, after probing we (as well as the optimum adversary) may choose *not to* include an item in the Knapsack, *arbitrarily (re)order* the schedule, come up widely different clustering depending on the outcome of the probed values. Hence, we need completely different arguments and observation schemes in the absence of the irrevocable commitment.

The classic stochastic optimization (non-adaptive, non-probing) versions of these scheduling problems were considered in [28, 16]. We also note that stochastic optimization problems were considered in [26, 23, 25, 34, 36, 22, 24]; these problems appear to be unrelated to adaptivity gaps.

2 General Framework

The types of objectives we are interested in this paper are scheduling and metric problems. For these problems, the key feature is that the final solution has to be constructed over *all* the variables whether probed or unprobed. Our main contribution is to identify a key **recombinant** property and an uniform solution recipe that shows constant factor adaptivity gap for all problems which

¹We show in Section 3 that probing does indeed improve the objective of $1/\sum_j w_j C_j$ by a polynomial amount.

satisfy the property. We present this general technique below, and adapt it to specific problems in the subsequent sections.

2.1 Problems Considered

Before presenting the general algorithmic framework and analysis using the recombinant property, we summarize our problems and results.

Scheduling Problems. For scheduling we consider the average completion time ($1||\sum_j w_j C_j$) and minimum makespan on identical machines; in both cases, the job sizes are random variables. We focus on the weighted completion time which illustrates the main issues (Section 3). A surprising feature of this algorithm is that *only the expected values* of job sizes are used in constructing the approximately optimal probing scheme. The makespan problem is discussed in Section 4. For both these problems we design non-adaptive algorithm which shows that the adaptivity gap is a constant.

Metric Problems. We next consider several optimization problems in metric spaces. We assume that the input nodes are discrete distributions of polynomial specification over points of the metric space. For these problem a small but important added twist is needed – we need to reformulate the problem on a different but related metric. We consider the k -Median and Steiner Tree (or TSP) problem. We discuss the k -Median problem and the general setup for these problems in Section 5. The MST problem is discussed in Section 6. For both these problems we design non-adaptive algorithm which shows that the adaptivity gap is a constant.

For the k -median problem an interesting issue comes to fore, which does not appear for deterministic input. It is well known that over any space, there exists a k -median solution which uses the input points and is at most twice the optimum. For distributional input we show that the adaptivity gap for obtaining exactly k medians is polynomially large, short of a polynomial blowup in the probing cost. However, if we restrict all solutions to use fixed points in the metric space as medians (as opposed to declaring an input node which could possibly be a distribution over points as a median) then the gap disappears! This exposes an interesting contrast in the problem based on which points are allowed to be medians. We expect these definitional issues to be of independent interest as more problems with distributions as input are investigated.

2.2 The Non-adaptive Algorithm

For $S \subseteq [n]$, let $\mathcal{O}_S$ denote the space of solutions constructed on the subset S of input variables. Let $\mathcal{O} = \mathcal{O}_{[n]}$. For instance, in weighted completion time scheduling, $\mathcal{O}_S$ denotes the space of orderings of the jobs in subset S . Given a realization $\mathbf{v}_A$ of variables $\{X_i, i \in A\}$, let $\min_{o \in \mathcal{O}_A} h_A(o, \mathbf{v}_A)$ denote the optimal objective using only these realized variables.

Definition 2.1. *The outlier problem $\mathcal{G}^*(C)$ is: Choose a subset S of variables so that $\sum_{i \in S} c_i \leq C$ as outliers (i.e., to ignore), so that $\min_{o \in \mathcal{O}_{\bar{S}}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h_{\bar{S}}(o; \mathbf{v}_{\bar{S}})]$ is minimized. Here, $\bar{S} = [n] \setminus S$.*

As an example, in weighted completion time scheduling, $\mathcal{G}^*(C)$ would correspond to choosing a subset S of jobs with total probing cost at most C as outliers (i.e., to ignore) so that for the remaining jobs $\bar{S}$, the expected completion time (with no probing), $\sum_{i,j \in \bar{S}, j \geq i} w_j \mu_i$ is minimized. Here, $\mu_i = \mathbf{E}[X_i]$.

Our non-adaptive strategy has the following simple structure:

1. Solve the *outlier* problem $\mathcal{G}^*(C)$. Suppose this outputs a subset $S \subseteq [n]$ of as outliers.

2. The non-adaptive algorithm probes S , and for the realization $\mathbf{v}_S$ of these variables, subsequently solves $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$ on the observed and unobserved variables.

As an example, in weighted completion time scheduling, suppose the outlier problem returns a subset S of jobs as outliers. The probing strategy observes the sizes of these jobs. It then sorts the jobs in decreasing order of $\frac{w_i}{l_i}$ and schedules them, where $l_i = \mu_i$ if $i \in \bar{S}$, and $l_i = \mathbf{v}_i$ if $i \in S$.

2.3 Analysis

The guarantees we obtain will depend on the approximation ratio achievable for $\mathcal{G}^*(C)$ and for solving $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$. We will require two properties about the problems we consider. The first property is *downward closure*.

$$\forall o \in \mathcal{O}_{[n]}, A \subseteq [n], \quad \exists o_A \in \mathcal{O}_A \text{ s.t.} \quad \forall \mathbf{v}_{[n]} \quad h_A(o_A; \mathbf{v}_A) \leq h(o; \mathbf{v}_{[n]}) \quad (\text{P1})$$

The above property trivially holds for all the problems we consider: weighted completion time scheduling and minimum makespan scheduling where $\mathbf{v}$ denotes job sizes; and geometric k -medians and Steiner trees, where $\mathbf{v}$ denotes locations of points in a metric space.

Using (P1), we will show a lower bound for the adaptive optimal solution in terms of the outlier problem. In particular, we relate the quantity $\mathcal{G}^*(C)$ to the values OPT_s and OPT_h of the optimal adaptive solutions with soft and hard probing budgets C respectively. Recall that in the hard budget version, the probe cost has to be at most C on all adaptive decision paths, whereas in the soft budget version, it is only required that the expected probe cost be at most C . Clearly, $OPT_h \geq OPT_s$ since the soft budget version relaxes the hard budget constraint.

Lemma 2.2. $\mathcal{G}^*(C) \leq OPT_h$, and $\mathcal{G}^*((1 + \beta)C) \leq \left(1 + \frac{1}{\beta}\right) OPT_s$ for all $\beta > 0$.

Proof. First consider the optimal decision tree with hard budgets. At each leaf l of this tree, the probing cost is at most C . Let S denote the subset of probed variables corresponding to this leaf node, and $\mathbf{v}_S$ denote the values observed for these variables. Let opt_l denote the objective function at this leaf node. By property (P1), we have:

$$opt_l = \min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})] \geq \min_{o_{\bar{S}} \in \mathcal{O}_{\bar{S}}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o_{\bar{S}}; \mathbf{v}_{\bar{S}})] \geq \mathcal{G}(C^*)$$

Therefore, $opt_l \geq \mathcal{G}^*(C)$. Since this is true for all leaves l , we have $OPT_h \geq \mathcal{G}^*(C)$.

Next consider the optimal decision tree with soft budget C . Consider all leaves l with probing cost at most $(1 + \beta)C$. By Markov's inequality, the probability of such a leaf is at least $\frac{\beta}{1 + \beta}$. For each of these leaves, $opt_l \geq \mathcal{G}^*((1 + \beta)C)$. Therefore using (P1), we have:

$$OPT_s \geq \frac{\beta}{1 + \beta} \mathcal{G}^*((1 + \beta)C) \quad \Rightarrow \quad \mathcal{G}^*((1 + \beta)C) \leq \left(1 + \frac{1}{\beta}\right) \cdot OPT_s$$

□

Recombinant Property. The next and more non-trivial property relates to combining the solutions for the probed and unprobed parts. This property crucially requires that the random variables X_i are independent, and holds for all product distributions. In the inequality below, the expectations are over $\mathbf{v}_{\bar{A}}$.

$$\begin{aligned}
& \forall A \subseteq [n], \mathbf{v}_A, \min_{o \in \mathcal{O}} \mathbf{E}[h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})] \\
& \leq \rho \left(\min_{o \in \mathcal{O}_A} h_A(o; \mathbf{v}_A) + \min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})] + \mathbf{E}[\min_{o \in \mathcal{O}} h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})] \right)
\end{aligned} \tag{P2}$$

At a high level, the recombinant property holds if for an arbitrary partitioning of the input variables, each of the parts have an induced solution with the property that the objective function values sum to no more than the respective quantities in the original problem; further, the two solutions can be combined without a significant increase in value of the resulting solution. Since encoding the value of the probed part is unwieldy, this partitioning is necessary to construct a solution that encodes just the unprobed part. The recombination shows a small adaptivity gap – the tricky part is to ensure that the interaction of the probed and unprobed parts can be bounded. This technique also yields non-adaptive probing strategies in addition to the adaptivity gap proof.

For all the problems we consider, we will show that ρ is a constant. Using this, we can now derive the approximation ratio for the non-adaptive strategy. Suppose Step (1), the outlier problem, has approximation ratio γ_1 , and Step (2), solving $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}}[h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$ given $\mathbf{v}_S$ has approximation ratio γ_2 , then we have the following result:

Theorem 2.3. *The non-adaptive algorithm finds a solution of value at most $\rho\gamma_1(2 + \gamma_2)OPT_h$. It further finds a solution of value $\rho\gamma_1(2 + \gamma_2) \left(1 + \frac{1}{\beta}\right) OPT_s$ using outlier cost $(1 + \beta)C$ for any $\beta > 0$.*

Proof. Given the computed set of outliers S and the final solution $\nu_{\mathbf{v}_S}$ for the hard budget case, we have the following sequence of inequalities:

$$\begin{aligned}
& \mathbf{E}_{\mathbf{v}_S} [\mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(\nu_{\mathbf{v}_S}; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]] \\
& \leq \gamma_2 \mathbf{E}_{\mathbf{v}_S} \left[\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})] \right] \\
& \leq \gamma_2 \rho \mathbf{E}_{\mathbf{v}_S} \left[\min_{o \in \mathcal{O}_S} h_A(o; \mathbf{v}_S) + \min_{o \in \mathcal{O}_{\bar{S}}} \mathbf{E}[h_{\bar{S}}(o; \mathbf{v}_{\bar{S}})] + \mathbf{E}[\min_{o \in \mathcal{O}} h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})] \right] \\
& \leq 2\gamma_2 \rho \mathbf{E}_{\mathbf{v}} \left[\min_{o \in \mathcal{O}} h(o; \mathbf{v}) \right] + \gamma_2 \rho \mathbf{E}_{\mathbf{v}_S} \left[\min_{o \in \mathcal{O}_{\bar{S}}} \mathbf{E}[h_{\bar{S}}(o; \mathbf{v}_{\bar{S}})] \right] \\
& \leq 2\gamma_2 \rho \mathbf{E}_{\mathbf{v}} \left[\min_{o \in \mathcal{O}} h(o; \mathbf{v}) \right] + \gamma_2 \rho \gamma_1 \mathcal{G}^*(C) \\
& \leq \gamma_2 \rho (\gamma_1 + 2) OPT_h
\end{aligned}$$

The second inequality follows from Property (P2), the third from Property (P1), and the final inequality from Lemma 2.2. The proof for the soft budget case is identical. $\square$

Definition 2.4. *We will denote $\eta = \gamma_2 \rho (\gamma_1 + 2)$ as the approximation ratio of the problem.*

The above guarantees imply bicriteria approximations (where the probing cost also increases by a constant factor) for the soft budget versions of all problems, and for the hard budget versions of the metric problems we consider. The increase in probing cost is unavoidable using our techniques, since the best known algorithms for the outlier versions of the metric problems we consider have similar gaps. An interesting open question is to show a complexity result that the increase in budget is unavoidable.

3 Scheduling I: Weighted Completion Time on Single Machine

We first consider the weighted completion time problem ($1||\sum w_j C_j$) of scheduling jobs on a single processor to minimize the sum of the weighted completion times. All jobs are released at time $t = 0$ and there are no deadlines or precedence constraints.

In the probing model the sizes (or processing times) of jobs $J_1, \dots, J_n$ are distributed according to independent random variables $X_1, X_2, \dots, X_n$ respectively. The weight of job J_i is w_i , which is *not* a random variable. Let $\mathbf{E}[X_i] = \mu_i$. Each variable X_i corresponding to the size of job J_i has probing cost c_i ; probing yields its exact value. Let C denote the (soft) budget on probing cost.

The solution is a strategy for adaptively probing a subset the jobs so that the expected weighted completion time of scheduling *all* the jobs after the outcome of the probes is known, is minimized. This expectation is over the outcome of the probes, and over the distribution of the processing times of the unprobed jobs. We note that the scheduling policy fixes the ordering $o \in \mathcal{O}$ of all jobs after the results of the probes (say set S) are known, but before the sizes of the unprobed jobs are revealed. Therefore, the optimal scheduling policy whose value is $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$ will simply order the jobs in decreasing order of weight to the ratio of (expected) processing time (*Smith's Rule*) – this processing time is exactly known for probed jobs S , and is the expected processing time for unprobed jobs $\bar{S}$ ².

Benefit of Probing. To gain intuition, we present a simple example where probing helps by a factor of $\Omega(n)$. There are n jobs with unit weights and unit probing costs, and sizes which are 0 w.p. $1 - 1/n$, and 1 with the remaining probability. If no jobs are probed, each job has expected size $1/n$, so that the expected completion time of any ordering is $\sum_{i=1}^n i/n = \frac{n+1}{2}$. If all jobs are probed, the jobs which are of size 1 can be placed after jobs of size 0. If there are k jobs of size 1, the completion time is $\frac{k(k+1)}{2}$. The variable k follows Binomial($n, 1/n$), with mean 1 and variance at most 1. This implies $\mathbf{E}[k^2] \leq 2$, so that $\mathbf{E} \left[\frac{k(k+1)}{2} \right] \leq \frac{3}{2}$. Therefore, probing yields a $\Omega(n)$ benefit.

Main Result. Recall that ρ is the approximation for Property (P2), γ_1 is the approximation ratio for the outlier problem $\mathcal{G}(C^*)$, and γ_2 is the approximation ratio for solving $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$. We will show the following theorem.

Theorem 3.1. *For the weighted completion time scheduling problem, $\rho = 2$, $\gamma_1 = 3$ using cost budget $3C$, and $\gamma_2 = 1$. This implies an approximation ratio of $\eta = 10$ for the non-adaptive probing strategy that uses cost budget $3C$.*

The proof that $\gamma_2 = 1$ is simple, since the optimal scheduling policy whose value is $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$ will simply order the jobs in decreasing order of the ratio of weight to (expected) processing time (*Smith's Rule*) – this processing time is exactly known for probed jobs S , and is the expected processing time for unprobed jobs $\bar{S}$. We will first show $\rho = 2$, and then design a 3-approximation for the outlier problem that violates the cost budget by a factor of 3.

3.1 The Recombinant Property (P2)

We first show Property (P2) holds for $\rho = 2$. Consider n jobs with *deterministic* lengths. Let $l_1, l_2, \dots, l_n$ denote the job-lengths, and $w_1, w_2, \dots, w_n$ denote the job weights. Let $\alpha_i = \frac{l_i}{w_i}$. By Smith's rule, the optimal solution sorts the jobs in increasing order of α_j and schedules in this order.

²This is verified by estimating the benefit/loss of an exchange of consecutive jobs on each scenario and then aggregating over the scenarios, see also [32].

The completion time of the optimal ordering can therefore be written as $\sum_{i=1}^n \sum_{j \geq i} w_i w_j \min(\alpha_i, \alpha_j)$. We have:

Lemma 3.2. *For any partitioning of n deterministic jobs into two disjoint sets A and B , we have:*

$$\sum_{i,j \in A, j \geq i} w_i w_j \min(\alpha_i, \alpha_j) + \sum_{i,j \in B, j \geq i} w_i w_j \min(\alpha_i, \alpha_j) \geq \sum_{i \in A, j \in B} w_i w_j \min(\alpha_i, \alpha_j)$$

Proof. Let $\gamma_{ij} = \min(\alpha_i, \alpha_j)$. First, suppose all $\gamma_{ij} = 1$. We have:

$$\begin{aligned} & \left(\sum_{i \in A} w_i - \sum_{j \in B} w_j \right)^2 \geq 0 \\ \Rightarrow & 2 \left(\sum_{i,j \in A, j \geq i} w_i w_j + \sum_{i,j \in B, j \geq i} w_i w_j \right) \geq 2 \left(\sum_{i \in A} w_i \right) \left(\sum_{j \in B} w_j \right) \\ \Rightarrow & \sum_{i,j \in A, j \geq i} w_i w_j + \sum_{i,j \in B, j \geq i} w_i w_j \geq \sum_{i \in A, j \in B} w_i w_j \end{aligned}$$

We next consider the case of general α_i . We will prove this by induction on the number of jobs (the base case being trivial). Let $i^* = \operatorname{argmin}_i \alpha_i$. For each job i , let $\beta_i = \alpha_i - \alpha_{i^*}$, and let $\delta_{ij} = \min(\beta_i, \beta_j)$. We have:

$$w_i w_j \gamma_{ij} = w_i w_j (\alpha_{i^*} + \delta_{ij})$$

From the proof of the $\gamma_{ij} = 1$ case, we have:

$$\alpha_{i^*} \left(\sum_{i,j \in A, j \geq i} w_i w_j + \sum_{i,j \in B, j \geq i} w_i w_j \right) \geq \alpha_{i^*} \sum_{i \in A, j \in B} w_i w_j$$

The set of jobs with non-zero β values is strictly smaller than n . Let Z be the set of jobs with $\beta_i = 0$. By the inductive hypothesis we have:

$$\sum_{i,j \in A \setminus Z, j \geq i} w_i w_j \delta_{ij} + \sum_{i,j \in B \setminus Z, j \geq i} w_i w_j \delta_{ij} \geq \sum_{i \in A \setminus Z, j \in B \setminus Z} w_i w_j \delta_{ij}$$

Adding the previous two inequalities, we have the proof of the lemma. $\square$

In the above lemma, the LHS represents the contribution to the optimal completion time which arises from job pairs *within* A and within B . The RHS represents contributions of job pairs such that one of the jobs is in A and the other in B . The above shows that the interaction term across the two sides of any partition, can be bounded by the sum of the interactions within each side. This directly shows the recombinant property (P2) with $\rho = 2$:

Lemma 3.3.

$$\forall A \subseteq [n], \mathbf{v}_A, \quad \min_{o \in \mathcal{O}} \mathbf{E}[h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})] \leq 2 \left[\min_{o \in \mathcal{O}_A} h_A(o; \mathbf{v}_A) + \min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})] \right]$$

where all expectations are over $\mathbf{v}_{\bar{A}}$.

Proof. Let $\mu_i = \mathbf{E}[X_i]$. We have:

$$\begin{aligned}\min_{o \in \mathcal{O}_A} h_A(o; \mathbf{v}_A) &= \sum_{i,j \in A, j \geq i} \min(w_j \mathbf{v}_i, w_i \mathbf{v}_j) \\ \min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})] &= \sum_{i,j \in \bar{A}, j \geq i} \min(w_j \mu_i, w_i \mu_j)\end{aligned}$$

Furthermore, we have:

$$\begin{aligned}\min_{o \in \mathcal{O}} \mathbf{E}[h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})] &= \sum_{i,j \in A, j \geq i} \min(w_j \mathbf{v}_i, w_i \mathbf{v}_j) + \sum_{i,j \in \bar{A}, j \geq i} \min(w_j \mu_i, w_i \mu_j) + \sum_{i \in A, j \in \bar{A}} \min(w_j \mathbf{v}_i, w_i \mu_j) \\ &= \min_{o \in \mathcal{O}_A} h_A(o; \mathbf{v}_A) + \min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})] + \sum_{i \in A, j \in \bar{A}} \min(w_j \mathbf{v}_i, w_i \mu_j)\end{aligned}$$

By Lemma 3.2, we have:

$$\sum_{i \in A, j \in \bar{A}} \min(w_j \mathbf{v}_i, w_i \mu_j) \leq \sum_{i,j \in A, j \geq i} \min(w_j \mathbf{v}_i, w_i \mathbf{v}_j) + \sum_{i,j \in \bar{A}, j \geq i} \min(w_j \mu_i, w_i \mu_j)$$

Putting the above two inequalities together completes the proof. $\square$

3.2 Approximation Algorithm for the Outlier Problem

To complete the proof of Theorem 3.1, we finally focus on solving the outlier problem $\mathcal{G}^*(C)$. Consider the following integer program for the outlier program. Here, variable $z_i \in \{0, 1\}$ is set to 1 if job i is in the outlier set. Variable $e_{ij} \in \{0, 1\}$ is set to 1 if jobs i and j are both not in the outlier set.

$$\begin{aligned}\min \quad & \sum_{i=1}^n \sum_{j>i} e_{ij} \min\{w_j \mu_i, w_i \mu_j\} + \sum_{i=1}^n (1 - z_i) w_i \mu_i \\ & \sum_{i=1}^n z_i c_i \leq C \\ & e_{ij} + z_i + z_j \geq 1 \quad \text{for all } i, j > i \\ & e_{ij}, z_i \in \{0, 1\} \quad \text{for all } i, j \in \{1, 2, \dots, n\}\end{aligned}$$

It is easy to check that this integer program encodes the problem $\mathcal{G}^*(C)$. To see this, note that the objective is precisely the weighted completion time of the jobs not in the outlier set. The first constraint encodes the cost constraint on the outlier set, and the second constraint encodes that if jobs i and j are both not in the outlier set, then $e_{ij} = 1$, but if z_i or z_j is 1, then $e_{ij} = 0$.

We solve the linear relaxation of the LP, where the final constraint is replaced with $e(i, j) \geq 0$ and $z_i \geq 0$. Now round the LP solution as follows: Let $S = \{i | z_i \geq \frac{1}{3}\}$. Set $\tilde{z}_i = 1$ if $i \in S$, and set $\tilde{e}_{ij} = 1$ if $i \notin S$ and $j \notin S$. The set S is the outlier set output by the algorithm.

Theorem 3.4. *The outlier problem $\mathcal{G}^*(C)$ admits to a 3-approximation that violates the cost budget C by a factor of 3.*

Proof. It is easy to check that $\tilde{z}_i \leq 3z_i$ and $\tilde{e}_{ij} \leq 3e_{ij}$. To see this, simply observe that $e_{ij} = 1$ iff $z_i < 1/3$ and $z_j < 1/3$, in which case $e_{ij} \geq 1/3$. Therefore, the objective is within a factor 3 of the LP objective, and the first constraint is violated by a factor of 3. $\square$

4 Scheduling II: MAKESPAN on Identical Machines

We now consider the problem of minimizing the makespan on identical parallel machines. In this problem, there are m identical machines, and n jobs, whose sizes are random variables, $X_1, X_2, \dots, X_n$. We can probe job i by spending cost c_i , and find the exact value of its processing time. Given a bound on the total query cost C , the goal is to find the subset of variables to probe so that the expected value of the makespan (*i.e.* the load on the most loaded machine) is minimized. Here, the expectation is over all realizations of the sizes of the unprobed jobs.

Recall that ρ is the approximation for Property (P2), γ_1 is the approximation ratio for the outlier problem $\mathcal{G}(C^*)$, and γ_2 is the approximation ratio for solving $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$. We will show the following theorem.

Theorem 4.1. *For the minimum makespan scheduling problem, we have $\gamma_1 = O(1)$, $\gamma_2 = O(1)$, and $\rho = 1$, so that the approximation ratio of the non-adaptive strategy is $\eta = O(1)$.*

Recombinant Property. It is easy to see that for any $A \in [n]$, for all $o_1 \in \mathcal{O}_A$ and $o_2 \in \mathcal{O}_{\bar{A}}$, the concatenation of these solutions (call it $o \in \mathcal{O}$) satisfies:

$$h(o; \mathbf{v}) \leq h(o_1; \mathbf{v}_A) + h(o_2; \mathbf{v}_{\bar{A}}) \quad \forall \mathbf{v}$$

This directly implies Property (P2) with $\rho = 1$.

Outlier Problem. The outlier version $\mathcal{G}^*(C)$ is the following: Find a subset S of jobs with cost at most C to discard so that the optimal expected makespan on the remaining jobs is minimized. We therefore consider the problem $\mathcal{P}$ of scheduling the jobs in the absence of probing. The goal is simply to assign the jobs to the machines so that the expected makespan, which is the expected load of the most loaded machine, is minimized. Here, the expectation is over all possible realizations of the job sizes. This problem was addressed by Kleinberg, Rabani, and Tardos [28], who show a nice characterization.

Definition 4.2. *Given a random variable X_i , define the random variable $Y_{i,t}, Z_{i,t}$ as follows*

$$Y_{i,t} = \begin{cases} X_i/t & \text{if } X_i \leq t \\ 0 & \text{otherwise} \end{cases} \quad Z_{i,t} = \begin{cases} 0 & \text{if } X_i \leq t \\ X_i & \text{otherwise} \end{cases}$$

In other words Y is the contribution of X below a threshold t and Z is the contribution of X above t ; naturally $X_i = t \cdot Y_{i,t} + Z_{i,t}$. Let

$$f_i(t) = \sum_i \left(E[Z_{i,t}] + \frac{t \log E[m^{Y_{i,t}}]}{\log m} \right)$$

Theorem 4.3 ([28]). *Given a set of jobs whose sizes follow independent random variables $\{X_i\}$,*

- *If $\sum_i f_i(t) \geq t/3$ then any scheduling of the jobs on m machines has expected makespan $\Omega(t)$.*
- *If $\sum_i f_i(t) \leq 2t/3$ then the following algorithm produces a schedule with (expected) makespan $O(t)$:*

1. *Define the “effective size” of job i to be $\eta_i(t) = \frac{\log E[m^{Y_{i,t}}]}{\log m}$.*
2. *Consider the jobs in arbitrary order, and place each job on that machine where the sum of the $\eta_j(t)$ of the already scheduled jobs j is minimum. In other words, use Graham’s rule on $\eta_j(t)$.*

The following algorithm solves $\mathcal{G}^*(C)$ to a $O(1)$. Try different values of t in increasing powers of $(1 + \epsilon)$. For each t , decide if there is a subset S of jobs so that: (1) $\sum_{i \in S} c_i \leq C$, and (2) $\sum_{i \notin S} f_i(t) \leq t/2$. By Theorem 4.3, the smallest such t (call it t^*) for which the decision problem returns a “yes” answer is the desired $O(1)$ approximation to $\mathcal{G}^*(C)$. It is easy to see that t^* will be at most the sum of the maximum values the X_i can take. Therefore, since the distributions are discrete and specified as input, computing $f_i(t)$ takes polynomial time for any t of interest. This decision problem is a knapsack problem, and can be solved to a $(1 + \epsilon)$ approximation on t^* without violating the cost budget. This shows $\gamma_1 = O(1)$.

To show that $\gamma_2 = O(1)$, we simply schedule the probed jobs using Graham’s rule on the observed sizes. Separately schedule the unprobed jobs using the algorithm from [28], and append this schedule to that of the probed jobs. Since the probed jobs are scheduled to a factor 2 approximation, and the unprobed jobs are scheduled to a $O(1)$ approximation (by the result of [28]), this shows that $\gamma_2 = O(1)$, completing the proof of Theorem 4.1.

5 Metric Problems I: K -median Clustering

We next consider several metric problems, specifically the k -median clustering and the minimum spanning tree problem on metric spaces. Before defining the problems, we will define the uncertainty model.

5.1 Uncertainty Model for Metric Problems

We define the uncertainty model as follows. We are given a metric space with point set $\mathcal{P}$, which defines a distance function l . The input is a set of nodes $\mathcal{V}$, where the location of node i follows an independent distribution X_i over $\mathcal{P}$. Distribution (or node) $i \in \mathcal{V}$ has probe cost c_i ; on probing, the node resolves to one of the locations in $\mathcal{P}$. Since we can only probe a subset of nodes, but must finally construct a solution over all the nodes, it will be helpful to define a metric space over the set of points $\mathcal{P}$ and nodes $\mathcal{V}$, *i.e.*, over the set $\mathcal{M} = \mathcal{P} \cup \mathcal{V}$. For $i, j \in \mathcal{V}$, let $D(i, j) = l(X_i, X_j)$ denote the random variable corresponding to the distance between X_i and X_j . Let $d(i, j) = \mathbf{E}[D(i, j)]$ where the expectation is over the random variables X_i and X_j .

Claim 5.1. $d(i, j) + d(j, k) \geq d(i, k)$.

Proof. For any realization of the values of X_i, X_j , and X_k , we have $D(i, j) + D(j, k) \geq D(i, k)$. Taking expectations over the random choices of X_i, X_j , and X_k , the claim follows. $\square$

Define the following metric space: The vertices are the points $\mathcal{M} = \mathcal{P} \cup \mathcal{V}$. The distance metric is d . For $i, j \in \mathcal{V}$, $d(i, j) = \mathbf{E}[D(i, j)] = \mathbf{E}[l(X_i, X_j)]$. For $i, j \in \mathcal{P}$, $d(i, j) = l(i, j)$. For $i \in \mathcal{P}, j \in \mathcal{V}$, define $d(i, j) = \mathbf{E}[l(i, X_j)]$. Using the above claim, it is clear that the function d defines a metric space on the nodes $\mathcal{V} \cup \mathcal{P}$. This completes the modeling of uncertainty for the metric problems we consider.

5.2 K -medians: Problem Statement

In this problem, we are given a metric space with point set $\mathcal{P}$, which defines a distance function l . The input is a set of nodes $\mathcal{V}$, where the location of node i follows an independent distribution X_i over $\mathcal{P}$. Distribution $i \in \mathcal{V}$ has probe cost c_i . The goal is to design an adaptive policy to probe the nodes which spends expected cost at most C . After probing, the algorithm opens K centers and assigns all probed and unprobed nodes to some center so that the expected distance cost (or *value*)

of the clustering is minimized. This expectation is over the locations of the unprobed nodes. Note that after probing, the center selection and assignment policy assigns an unprobed node i to that open center w which minimizes the expected distance $\mathbf{E}[l(X_i, w)]$ where the expectation is over the random variable X_i . The goal is to design a probing policy whose resulting expected distance cost (or value) of K -median clustering is minimized, where the expectation is over the outcomes of the probes and the locations of the unprobed nodes.

We consider two variants of the problem. In the first variant, we assume that the center selection policy is restricted to opening centers from a set $S \subseteq \mathcal{P}$. This means the centers can only be chosen from points of the underlying metric space. Therefore, for an unprobed node i assigned to a center $w \in \mathcal{P}$, the expected distance cost is $\mathbf{E}[l(X_i, w)]$, where the expectation is over the random variable X_i . In the second variant, the centers are allowed to be *input nodes*, and therefore distributions. Now, an unprobed node j can be opened as a center after probing a set of nodes. Suppose an unprobed node i is assigned to this center, then the expected distance cost is $\mathbf{E}[l(X_i, X_j)]$, where the expectation is over *both* the random variables X_i and X_j .

We present a constant factor adaptivity gap for the former variant. We then show that the adaptivity gap is polynomially large for the latter variant. This shows a fundamental difference in the two variants.

5.3 Fixed Centers

We first consider the first variant where the centers can only be points from $S \subseteq \mathcal{P}$. Recall that ρ is the approximation for Property (P2), γ_1 is the approximation ratio for the outlier problem $\mathcal{G}(C^*)$, and γ_2 is the approximation ratio for solving $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$. We will show the following theorem.

Theorem 5.2. *For the K -median problem with fixed centers, $\rho = 5$, $\gamma_1 = 5$ with outlier cost $5C$, and $\gamma_2 = 3 + \epsilon$. This shows an approximation ratio of $\eta = O(1)$ with outlier cost $5C$.*

As shown in Section 5.1, first define the new metric space d over the points $\mathcal{P} \cup \mathcal{V}$. Given the set of probed values $\mathbf{v}_S \subseteq \mathcal{P}$, and the unprobed nodes $\bar{S} \subseteq \mathcal{V}$, the problem reduces to deterministic K -medians on $\mathbf{v}_S \cup \bar{S} \subseteq \mathcal{P} \cup \mathcal{V}$ in metric space d , and this shows $\gamma_2 = 3 + \epsilon$ using the algorithm in [2]. In the outlier problem, the goal is to find the subset $T \subseteq \mathcal{V}$ of nodes of total probing cost at most C such that the cost of K -median clustering of the remaining nodes $\mathcal{V} \setminus T$ in metric space d is minimized. This problem has a $\gamma_1 = 5$ approximation due to the authors of [6] if the outlier cost is relaxed to $5C$. If all probing costs are unit, we have $\gamma_1 = O(1)$ approximation that does not relax the outlier cost [7]. The only part left to show is the recombinant property (P2).

Lemma 5.3.

$$\forall A \subseteq [n], \mathbf{v}_A, \min_{o \in \mathcal{O}} \mathbf{E}[h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})] \leq 5 \cdot \min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})] + 4 \cdot \mathbf{E}[\min_{o \in \mathcal{O}} h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})]$$

where the expectations are over $\mathbf{v}_{\bar{A}}$.

Proof. The quantity $Q_1 = \min_{o \in \mathcal{O}} \mathbf{E}[h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})]$ on the LHS involves constructing the optimal K -median solution using the actual realization $\mathbf{v}_A$ of the probed set A , and the expected locations of the unprobed nodes $\bar{A}$. To bound this cost, we first construct the optimal K medians on the unprobed nodes $\bar{A}$ by paying $Q_2 = \min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})]$. Next, move the unprobed nodes to the assigned median. For $i \in \bar{A}$, let $\phi(i)$ denote the assigned median.

We will now proceed by considering scenarios $\mathbf{v}_A, \mathbf{v}_{\bar{A}}$ of the values of all nodes. Fix some such scenario σ . Let M_σ denote the value of the optimal K -median solution in this scenario. The

expected value of this optimal solution over the realizations of $\mathbf{v}_{\bar{A}}$ is $Q_3 = \mathbf{E}[\min_{o \in \mathcal{O}} h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})]$. Let $P' \subseteq \mathcal{P}$ denote the set of points $\mathbf{v}_A$, along with the nodes $i \in \bar{A}$ located at point $\phi(i)$. Note that the set P_σ is the same for all realized σ . Since $\mathbf{v}_A$ is fixed, and the mapping ϕ is independent of the scenario, we have P' is independent of the scenario σ .

We will now construct the solution corresponding to the quantity Q_1 using the solution for Q_2 and the solutions M_σ . In scenario σ , send each point in A its assigned center in the optimal solution M_σ . Similarly, send each node $i \in \bar{A}$ located at $\phi(i)$ back to its realized location in scenario σ and from there to its assigned center in M_σ . This yields a mapping from the nodes P' to K medians in scenario σ . The expected distance value (over scenarios σ) of this mapping is at most $Q_2 + Q_3$ by triangle inequality. Note that the distances in this mapping are distances between points in $\mathcal{P}$, and do not involve the distributional nodes $\mathcal{V}$. This yields a valid K -median solution on P' .

Since there is a feasible K -median solution for each σ , these when averaged over σ define a fractional K -median solution for the set of points P' of value at most $Q_2 + Q_3$. Therefore, there is an integer K -median solution on P' of value 4 times this fractional value [5]. The final solution corresponding to Q_1 maps the points $i \in \bar{A}$ first to $\phi(i)$ paying cost Q_2 , and then maps all the points to these constructed centers paying cost $4(Q_2 + Q_3)$. This shows $Q_1 \leq 5Q_2 + 4Q_3$, completing the proof. $\square$

5.4 Arbitrary Centers

Consider now the case where the centers themselves are allowed to be input nodes, and therefore distributions. After probing, the center selection policy could decide to open an unprobed input node as a center, and assign probed and unprobed nodes to this center. The distance cost between the center and the assigned node is the expected distance between them, where the expectation is taken over possible locations of the center and the assigned node. We note that the linear programming relaxation is still a lower bound on the adaptive optimal solution.

By constructing separate K -median solutions for the probed and unprobed variables, it is easy to show a $O(1)$ approximation that uses $2K$ centers and pays probing cost $5C$. We show that this is the best possible result for this case in the following sense. We show that any non-adaptive algorithm has polynomially large adaptivity gap on *both* distance and probing cost if it is restricted to opening at most $(1 + \epsilon)K$ centers for some constant $\epsilon > 0$. Intuitively what fails in reducing the $2K$ centers to K centers is the following: In the proof for fixed centers, we mapped the unprobed nodes of the non-adaptive solution to a set of nodes in $\mathcal{P}$, and constructed a fractional solution using these locations for these nodes. This ensures that the metric space is over $\mathcal{P}$ and independent of distributions. In the current setting, an unprobed node could be mapped to an unprobed center. Therefore across scenarios, the metric space over the locations of the nodes changes.

Theorem 5.4. *The adaptivity gap for K -medians when centers can be input nodes is polynomially large on both distance and probing cost.*

Proof. Consider M distinct copies (at a mutual distance of at least M^3L from each other, $L = M^2$) of the following 2-dimensional node set. In copy m , there are $r + 1$ “cheap” nodes $X_{1m}, X_{2m}, \dots$ which cost 1 to probe. Distribution X_{im} is $(0, 0)$ with probability $1/2$ and $(i + 1, 0)$ otherwise. In addition, there are $t \gg \sqrt{M}$ pairs of nodes which are well-separated from other pairs by a large distance L . Pair j corresponds to two distributions: Y_{jm} and Z_{jm} . Y_{jm} is $(L + jL, 1)$ with probability $1 - \log t/t$ and $(L + jL, 0)$ with probability $\log t/t$. Z_{jm} is $(L + jL, -1)$ with probability $1 - \log t/t$ and $(L + jL, 0)$ with probability $\log t/t$. These distributions are “expensive” with probing cost $(r + 1)M$, where $r > 2 \log Mt$. Again note that the nodes for each m are far removed from the nodes corresponding to other m .

Let $K = (2t + r)M$. For each m , the adaptive solution will place $2t + r$ medians using the following strategy. First probe all cheap distributions X_* . There are two cases:

Case 1. For some m , X_{*m} resolve to $r + 1$ values distinct points. This happens with probability $1/2^{r+1}$ for a particular m and therefore $\frac{M}{2^{r+1}}$ overall. In this case, probe *all* the “expensive” distributions Y_* and Z_* . Choose the $r + 1$ realized points X_* , and some $2t - 1$ of the remaining $2t$ points as medians. With probability $1 - (1 - \log t/t)^{2t} \geq 1 - 1/t^2$ some (Y, Z) pair collides on the x -axis. In this case, the k medians solution has distance cost 0, else it has a distance cost of at most 2. The expected distance cost is therefore at most $(1/2^{r+1}) \cdot (1/t^2) \cdot 2$ for each m . Therefore, the overall expected distance cost is $\frac{M}{t^2 2^r}$.

Case 2. If at most r distinct values of X_{*m} are observed for all m , then do not probe further, since there is a k median solution of value 0, in which every one of the expensive nodes and all the realized locations of the cheap nodes are the medians.

The expected probing cost of this scheme is at most $M(r+1) + (r+1)M \cdot 2Mt \cdot \frac{M}{2^{r+1}} \leq 4M(r+1)$.

Any non-adaptive probing scheme must probe at least one expensive distribution in each copy, else the distance cost is at least $\frac{1}{2^{r+2}} \gg \frac{M}{t^2 2^r}$ in that copy: In the case where the cheap distributions resolve to distinct values w.p. $1/2^{r+1}$, the distance cost will be at least 0.5. Therefore, the probing cost needed is $(r+1)M^2$, which implies that unless the probing cost is a factor M larger than the adaptive scheme, the distance cost must be $\frac{t^2}{M}$ times larger. Therefore, no non-trivial adaptivity gap is possible. $\square$

6 Metric Problems II: Minimum Steiner Trees

We finally consider the minimum Steiner tree problem – the algorithm extends to Metric Traveling salesman problem naturally. As with K -medians, the input is a collection of n nodes. The location of node i is an independent random variable X_i , which is a distribution over points $\mathcal{P}$ in a metric space with distance function l . Let $\mathcal{V}$ denote the set of nodes. The exact location of node i is determined by spending probing cost c_i . The goal is to design an adaptive probing scheme which minimizes the expected cost of connecting the nodes by a spanning tree, subject to the constraint that this decision tree has expected probing cost at most C .

Recall that ρ is the approximation for Property (P2), γ_1 is the approximation ratio for the outlier problem $\mathcal{G}(C^*)$, and γ_2 is the approximation ratio for solving $\min_{o \in \mathcal{O}} \mathbf{E}_{\mathbf{v}_{\bar{S}}} [h(o; \mathbf{v}_S, \mathbf{v}_{\bar{S}})]$. We will show the following theorem.

Theorem 6.1. *For the metric Steiner tree problem, we have $\rho = 1$, $\gamma_1 = 4$ with outlier cost $4C$, and $\gamma_2 = 1.55$. The non-adaptive strategy therefore has an approximation ratio $\eta = O(1)$ with probing cost $4C$.*

Since the overall algorithm is similar to the K -medians problem, we simply highlight the differences. As discussed in Section 5.1, we construct the metric space d over $\mathcal{M} = \mathcal{P} \cup \mathcal{V}$. The outlier problem over the nodes $\mathcal{V}$ in the metric space d is defined as follows: There is a cost of c_i for every node $i \in \mathcal{V}$. The goal is to choose a set of nodes $S \subseteq \mathcal{V}$ with total cost at most C to discard such that the value of the Steiner tree on the nodes in $\mathcal{V} \setminus S$ is minimized. This has a $\gamma_2 = 4$ -approximation that spends outlier cost $4C$ [17]. If all probing costs are unit, we have $\gamma_2 = 4$ -approximation [14] that obeys the cost constraint. The minimum cost Steiner tree problem on the probed and unprobed nodes has an approximation ratio of $\gamma_2 = 1.55$ [33].

To show the recombinant property (P2) with $\rho = 1$, we make the following observations. For any set A of probed nodes with realization $\mathbf{v}_A$, $h_A(o; \mathbf{v}_A)$ is the value of the optimal Steiner tree

T_1 connecting these realized nodes; $\min_{o \in \mathcal{O}_{\bar{A}}} \mathbf{E}[h_{\bar{A}}(o; \mathbf{v}_{\bar{A}})]$ is the value optimal Steiner tree T_2 connecting the unprobed nodes; and finally, $\mathbf{E}[\min_{o \in \mathcal{O}} h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})]$ is at least the expected cost of any edge $e = (i, j)$ connecting some $i \in A$ and $j \in \bar{A}$. Taking the union of T_1 , T_2 , and e yields a tree whose cost is at least $\min_{o \in \mathcal{O}} \mathbf{E}[h(o; \mathbf{v}_A, \mathbf{v}_{\bar{A}})]$.

7 Conclusion

In this paper, we considered a large class of scheduling and metric problems when the inputs follow distributions that can be observed by paying a cost. We showed that the problem of computing the optimal budgeted resolution scheme for these inputs (that minimizes the expected value of the posterior optimization) is closely related to the outlier version of these problems. This work results in several open questions. First, can the approximation ratios be improved by an algorithm that designs adaptive resolution schemes instead of non-adaptive schemes? Next, our model for metric problems assumes nodes are uncertain and can be resolved at a cost. What if only *distances* between nodes can be resolved at a cost? The problem now is that the distributions for the edge lengths cannot be independent random variables, and we would need new techniques. Finally, it would be interesting to explore if the outlier scheme can be extended to a larger class of model-driven optimization problems.

Acknowledgments: We thank the anonymous referee for significantly improving the presentation of this paper. We also thank Sarel Har-Peled, Ashish Goel and Anupam Gupta for helpful discussions.

References

- [1] A. Akella, B. M. Maggs, S. Seshan, A. Shaikh, and R. K. Sitaraman. A measurement-based analysis of multihoming. In *ACM SIGCOMM Conference*, pages 353–364, 2003.
- [2] V. Arya, N. Garg, R. Khandekar, A. Meyerson, K. Munagala, and V. Pandit. Local search heuristics for k -median and facility location problems. *SIAM J. Comput.*, 33(3):544–562, 2004.
- [3] S. Babu and P. Bizarro. Proactive reoptimization. In *Proc. of the ACM SIGMOD Intl. Conf. on Management of Data*, 2005.
- [4] M. Charikar, R. Fagin, V. Guruswamy, J. Kleinberg, P. Raghavan, and A. Sahai. Query strategies for priced information. *Journal of Computer and System Sciences*, 64(4):785–819, 2002.
- [5] M. Charikar and S. Guha. Improved combinatorial algorithms for the facility location and k -median problems. In *FOCS '99: Proceedings of the 40th Annual Symposium on Foundations of Computer Science*, 1999.
- [6] M. Charikar, S. Khuller, D. M. Mount, and G. Narasimhan. Algorithms for facility location problems with outliers. In *SODA '01: Proceedings of the twelfth annual ACM-SIAM symposium on Discrete algorithms*, pages 642–651, 2001.
- [7] K. Chen. A constant factor approximation algorithm for k -median clustering with outliers. In *Proc. ACM-SIAM Symp. on Discrete Algorithms (SODA)*, 2008.
- [8] F. Chu, J. Halpern, and J. Gehrke. Least expected cost query optimization: What can we expect? In *Proc. of the ACM Symp. on Principles of Database Systems*, 2002.

- [9] F. Chu, J. Halpern, and P. Seshadri. Least expected cost query optimization: An exercise in utility. In *Proc. of the ACM Symp. on Principles of Database Systems*, 1999.
- [10] B. Dean, M. Goemans, and J. Vondrák. Approximating the stochastic knapsack problem: The benefit of adaptivity. *Math. Oper. Res.*, 33(4):945–964, 2008.
- [11] B. C. Dean, M. X. Goemans, and J. Vondrák. Adaptivity and approximation for stochastic packing problems. *Proceedings of SODA*, pages 395–404, 2005.
- [12] A. Deshpande, C. Guestrin, S. Madden, J. M. Hellerstein, and W. Hong. Model-driven data acquisition in sensor networks. In *Proc. of the 2004 Intl. Conf. on Very Large Data Bases*, 2004.
- [13] T. Feder, R. Motwani, R. Panigrahy, C. Olston, and J. Widom. Computing the median with uncertainty. *SIAM J. Comput.*, 32(2):538–547, 2003.
- [14] N. Garg. Saving an epsilon: A 2-approximation for the k -MST problem in graphs. In *STOC '05: Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, pages 396–402, 2005.
- [15] A. Goel, S. Guha, and K. Munagala. How to probe for an extreme value. *ACM Trans. Algorithms (to appear)*, 2008. Preliminary version appeared in *Proc. of the ACM Symp. on Principles of Database Systems (PODS)*, 2006.
- [16] A. Goel and P. Indyk. Stochastic load balancing and related problems. In *Proc. of the Annual Symp. on Foundations of Computer Science*, 1999.
- [17] M. X. Goemans and D. P. Williamson. A general approximation technique for constrained forest problems. *SIAM J. Comput.*, 24(2):296–317, 1995.
- [18] S. Guha and K. Munagala. Model-driven optimization using adaptive probes. In *Proc. ACM-SIAM Symp. on Discrete Algorithms*, pages 308–317, 2007.
- [19] S. Guha and K. Munagala. Sequential design of experiments via linear programming. *Computing Research Repository, arXiv:0805.2630*, 2008. Preliminary version appeared in the *ACM Symposium on Theory of Computing*, 2007.
- [20] S. Guha, K. Munagala, and S. Sarkar. Information acquisition and exploitation in multi-channel wireless systems. *Computing Research Repository, arXiv:0804.1724*, 2008. Preliminary version appeared in the *ACM SIGMETRICS Conference*, 2006.
- [21] P. K. Gummadi, H. V. Madhyastha, S. D. Gribble, H. M. Levy, and D. Wetherall. Improving the reliability of internet paths with one-hop source routing. In *6th Symposium on Operating System Design and Implementation (OSDI)*, pages 183–198, 2004.
- [22] A. Gupta and M. Pál. Stochastic steiner trees without a root. *Proc. of ICALP*, pages 1051–1063, 2005.
- [23] A. Gupta, M. Pál, R. Ravi, and A. Sinha. Boosted sampling: Approximation algorithms for stochastic optimization. In *Proc. of the Annual ACM Symp. on Theory of Computing*, 2004.
- [24] A. Gupta, M. Pál, R. Ravi, and A. Sinha. What about wednesday? Approximation algorithms for multistage stochastic optimization. *Proc. of APPROX-RANDOM*, pages 86–98, 2005.

- [25] A. Gupta, R. Ravi, and A. Sinha. LP rounding approximation algorithms for stochastic network design. volume 32, pages 345–364, 2007.
- [26] N. Immorlica, D. Karger, M. Minkoff, and V. Mirrokni. On the costs and benefits of procrastination: Approximation algorithms for stochastic combinatorial optimization problems. In *Proc. of the Annual ACM-SIAM Symp. on Discrete Algorithms*, 2004.
- [27] S. Khanna and W-C. Tan. On computing functions with uncertainty. In *Proc. of the ACM Symp. on Principles of Database Systems*, 2001.
- [28] J. Kleinberg, Y. Rabani, and É. Tardos. Allocating bandwidth for bursty connections. *SIAM J. Comput.*, 30(1):191–217, 2000.
- [29] A. Krause and C. Guestrin. Near-optimal nonmyopic value of information in graphical models. *Twenty-first Conference on Uncertainty in Artificial Intelligence (UAI 2005)*, pages 324–331, 2005.
- [30] R. H. Mohring, A. S. Schulz, and M. Uetz. Approximation in stochastic scheduling: the power of LP-based priority policies. *J. ACM*, 46(6):924–942, 1999.
- [31] C. Olston. *Approximate Replication*. PhD thesis, Stanford University, 2003.
- [32] M. Pinedo. *Sceduling: Theory Algorithms and Systems*. Prentice-Hall, 1995.
- [33] G. Robins and A. Zelikovsky. Improved steiner tree approximation in graphs. In *SODA '00: Proceedings of the eleventh annual ACM-SIAM symposium on Discrete algorithms*, pages 770–779, 2000.
- [34] D. Shmoys and C. Swamy. An approximation scheme for stochastic linear programming and its application to stochastic integer programs. *J. ACM*, 53(6):978–1012, 2006.
- [35] M. Skutella and M. Uetz. Stochastic machine scheduling with precedence constraints. *SIAM J. Comput.*, 34(4):788–802, 2005.
- [36] C. Swamy and D. B. Shmoys. Sampling-based approximation algorithms for multi-stage stochastic. In *FOCS '05: Proceedings of the 46th Annual IEEE Symposium on Foundations of Computer Science*, pages 357–366, 2005.